

INTERNATIONAL CENTRE FOR ECONOMIC RESEARCH



WORKING PAPER SERIES

Fabio Maccheroni and William H. Ruckle

BV AS A DUAL SPACE

Working Paper no. 29/2001
October 2001

**APPLIED MATHEMATICS
WORKING PAPER SERIES**



BV as a Dual Space*

Fabio Maccheroni[†]

Università Bocconi, Milano, Italia and
ICER, Torino, Italia

William H. Ruckle

Clemson University, Clemson, South Carolina, USA

October 2001

Abstract

Let \mathcal{C} be a field of subsets of a set I . It is well known that the space FA of all the finitely additive games of bounded variation on \mathcal{C} is the norm dual of the space of all simple functions on \mathcal{C} . In this paper we prove that the space BV of all the games of bounded variation on \mathcal{C} is the norm dual of the space of all simple games on \mathcal{C} . This result is equivalent to the compactness of the unit ball in BV with respect to the vague topology.

Key words: Set functions, duality, compactness, coalitional games.

1 Introduction

Let \mathcal{C} be a field of subsets of a set I . It is well known that the space FA of all the finitely additive games of bounded variation on \mathcal{C} , equipped with the total variation norm, is isometrically isomorphic to the norm dual of the space of all simple functions on \mathcal{C} , endowed with the sup norm (see [2] p. 258).

In this paper we establish a parallel result for the space BV of all the games of bounded variation on \mathcal{C} . To this end, we first introduce simple games on \mathcal{C} : the ones that assign non zero worth only to a finite number of elements of \mathcal{C} . Then we show that BV , equipped with the total variation norm, is isometrically isomorphic to the norm dual of the space of all simple games, endowed with a suitable norm. A first proof of this result builds on the compactness of the unit ball in BV with respect to the vague topology, proved by Marinacci [4].

Another contribution of this paper is showing that the duality of BV is indeed equivalent to the vague compactness of the unit ball. In order obtain this result, we provide a second *direct*

*We wish to thank Erio Castagnoli, Adriana Castaldo, an anonymous referee, and especially Massimo Marinacci for helpful discussions. Fabio Maccheroni gratefully acknowledges the financial support of MIUR and Università Bocconi.

[†]Corresponding Author: Istituto di Metodi Quantitativi, Università Bocconi, viale Isonzo 25, 20135 Milano, Italy. Phone: ++39 +02 5836 5606. Fax: ++39 +02 5836 5634. Email: fabio.maccheroni@uni-bocconi.it

proof of the dual nature of BV , and then observe that the vague topology can be viewed as the weak* topology induced by duality.

The paper is organized as follows. After a brief section of preliminaries, in Section 3 we use compactness to obtain duality. In Section 4, we go the other way around: we prove duality and obtain compactness as a corollary. Section 5 clarifies the connections between the two approaches.

2 Preliminaries

This section contains a few, well known, definitions and properties. We follow the notation of cooperative game theory, instead of the more common one of measure theory, since the spaces of set functions we consider are widely used in that field (see, e.g. [1]).

Let \mathcal{C} be a field of subsets of a non empty set I . A set function $v : \mathcal{C} \rightarrow \mathbb{R}$ is a *game* if $v(\emptyset) = 0$. A game on \mathcal{C} is *monotone* if $v(A) \leq v(B)$ whenever $A \subseteq B$.

A *chain* $\{S_i\}_{i=0}^n$ in \mathcal{C} is a finite strictly increasing sequence

$$\emptyset = S_0 \subset S_1 \subset \dots \subset S_{n-1} \subset S_n = I$$

of elements of \mathcal{C} . BV is the set of all games such that

$$\|u\| = \sup \left\{ \sum_{i=1}^n |u(S_i) - u(S_{i-1})| : \{S_i\}_{i=0}^n \text{ is a chain in } \mathcal{C} \right\} < \infty.$$

A game in BV is said to be of *bounded variation*. The pair $(BV, \|\cdot\|)$ is a Banach space, generated by all monotone games (see [1] pp. 14, 26-28).

Since BV is a (proper) subspace of $\mathbb{R}^{\mathcal{C}}$, it inherits a topology from the product topology of $\mathbb{R}^{\mathcal{C}}$. This is the weak topology generated by the projection functionals

$$\begin{aligned} p_A : BV &\rightarrow \mathbb{R} \\ u &\mapsto u(A) \end{aligned}$$

where $A \in \mathcal{C}$. A net $\{u_a\}$ converges to u in this topology iff $u_a(A) \rightarrow u(A)$ for all $A \in \mathcal{C}$ (we write $u_a \xrightarrow{\mathcal{C}} u$). This topology is called *vague topology* (for the analogy with the vague topology on the set of probability measures).

3 From compactness to duality

In this section, we start by observing that the unit ball in BV is compact in the vague topology, and we use this fact to show that BV is a dual space.

Theorem 1 (Marinacci) *The unit ball $U(BV) = \{u \in BV : \|u\| \leq 1\}$ is compact w.r.t. the vague topology.*

Proof. See [4]. The proof is based on the Tychonoff Theorem. □

Let X be the space of all the games which are non zero only on a finite number of elements of \mathcal{C} , the *simple games*. The pair (BV, X) is dual w.r.t. the following functional

$$\begin{aligned} D : BV \times X &\rightarrow \mathbb{R} \\ (u, x) &\mapsto \sum_{A \in \mathcal{C}} u(A) x(A) . \end{aligned}$$

That is:

- D is bilinear,
- if $D(u, x) = 0$ for all u , then $x = 0$,
- if $D(u, x) = 0$ for all x , then $u = 0$.

Therefore, X can be interpreted as a total subspace of the algebraic dual of BV by identifying $x \in X$ with the linear functional

$$\begin{aligned} D_x : BV &\rightarrow \mathbb{R} \\ u &\mapsto D(u, x) . \end{aligned}$$

For this reason, in the rest of this section, we will write $\langle u, x \rangle$ instead of $D(u, x)$. Some useful properties of X are stated in the following lemma, whose easy proof is omitted.

Lemma 2 X is a total subspace of the norm dual BV' of BV , and

$$\|x\|_{BV'} = \max_{\|u\|=1} \left(\sum_{A \in \mathcal{C}} u(A) x(A) \right)$$

for all $x \in X$.

Moreover, the topology $\sigma(BV, X)$ coincides with the vague topology.

Given a game u of bounded variation, let $J_u : X \rightarrow \mathbb{R}$ be the functional defined by

$$J_u(x) = \langle u, x \rangle$$

for all $x \in X$.

Theorem 3 Let X' be the norm dual of $(X, \|\cdot\|_{BV'})$. The operator

$$\begin{aligned} J : BV &\rightarrow X' \\ u &\mapsto J_u \end{aligned}$$

is an isometric isomorphism from BV onto X' . Moreover, J is a vague-weak* homeomorphism.

Proof. The operator J is well defined, linear, injective and $\|J\| \leq 1$. Let $U(BV)$ be the unit ball in BV , $U(X')$ be the unit ball in X' . For the Goldstine-Weston density Lemma $J(U(BV))$ is weak* dense in $U(X')$ (see, e.g., [3] p. 126).

Consider the vague topology on BV and the weak* topology on X' . Let $\{u_a\}$ be a net in BV . We have that $u_a \xrightarrow{\mathcal{C}} u$ iff $u_a \xrightarrow{\sigma(BV, X)} u$ iff $\langle u_a, x \rangle \rightarrow \langle u, x \rangle$ for all $x \in X$ iff $J_{u_a}(x) \rightarrow J_u(x)$ for all $x \in X$ iff J_{u_a} weak* converges to J_u (briefly $J_{u_a} \xrightarrow{w^*} J_u$). Thus J is vague-weak* continuous. Therefore, $J(U(BV))$ is weak* compact in X' and

$$J(U(BV)) = U(X').$$

Since $\|J\| \leq 1$, then $\|J_u\| \leq \|u\|$, for all $u \in BV$. Suppose that there exists $u \in BV$ such that $\|J_u\| < \|u\|$. Then $u \neq 0$ and $0 < \left\| J_{\frac{u}{\|u\|}} \right\| = \rho < 1$, set $w = \frac{u}{\|u\|}$. Since $\left\| J_{\frac{w}{\rho}} \right\| = 1$, it follows that $J_{\frac{w}{\rho}} \in U(X')$, but $\frac{w}{\rho} \notin U(BV)$ and J is injective, which is absurd. Thus $\|J_u\| = \|u\|$, for all $u \in BV$ and J is an isometry.

If $x' \in X' - \{0\}$ then there exists $u \in U(BV)$ s.t. $J_u = \frac{x'}{\|x'\|}$, hence $J_{\|x'\|u} = x'$. That is J is surjective. Since, for every net $\{u_a\}$ in BV , $u_a \xrightarrow{\mathcal{C}} u$ iff $J_{u_a} \xrightarrow{w^*} J_u$, then J is a vague-weak* homeomorphism. \square

4 From duality to compactness

In this section the opposite approach is adopted, we directly prove that BV is a dual space and use this fact to provide an alternative proof that the unit ball in BV is compact w.r.t. the vague topology, thus obtaining the equivalence of the two results.

We define the game $e_A : \mathcal{C} \rightarrow \mathbb{R}$ by

$$e_A(B) = \begin{cases} 1 & \text{if } B = A \\ 0 & \text{otherwise} \end{cases}$$

for all $A \in \mathcal{C} - \{\emptyset\}$, and $e_\emptyset = 0$. Being $x = \sum_{A \in \mathcal{C}} x(A) e_A$ for all $x \in X$, we have $X = \langle e_A : A \in \mathcal{C} \rangle$.

For each chain $\Omega = \{S_i\}_{i=0}^n$ in \mathcal{C} , define a seminorm on X by

$$\|x\|_\Omega = \max_{0 \leq k \leq n} \left| \sum_{i=k}^n x(S_i) \right| \quad (1)$$

for all $x \in X$. Let $X_\Omega = \langle e_A : A \in \Omega \rangle$. If $x \in X_\Omega$, we say that x depends on the chain Ω .

For all $x \in X$, set

$$\|x\| = \inf \sum_{l=1}^L \|x_l\|_{\Omega_l}$$

where the inf is taken over all finite decompositions $x = \sum_{l=1}^L x_l$ in which x_l depends on the chain Ω_l and $\|\cdot\|_{\Omega_l}$ is defined as in (1) for all $l = 1, \dots, L$.

Lemma 4 *The function $\|\cdot\| : X \rightarrow \mathbb{R}$ is a norm on X .*

Proof. It is easy to prove that $\|\cdot\|$ is a seminorm. We just show that $x \neq 0$ implies $\|x\| \neq 0$. If $x \in X_\Omega$ and $A \in \Omega$ then

$$\|x\|_\Omega \geq \frac{1}{2} |x(A)|.$$

In fact,

$$|x(A)| = \left| \sum_{i:S_i \supseteq A} x(S_i) - \sum_{i:S_i \supset A} x(S_i) \right| \leq \left| \sum_{i:S_i \supseteq A} x(S_i) \right| + \left| \sum_{i:S_i \supset A} x(S_i) \right| \leq 2 \|x\|_\Omega$$

Let $x \in X - \{0\}$. There exists $A \in \mathcal{C}$ such that $x(A) \neq 0$. If $x = \sum_{l=1}^L x_l$, $x_l \in X_{\Omega_l}$ for all $l = 1, \dots, L$, then $A \in \Omega_l$ for some l (otherwise $x(A) = 0$). Hence,

$$|x(A)| = \left| \sum_{l:A \in \Omega_l} x_l(A) \right| \leq \sum_{l:A \in \Omega_l} |x_l(A)|,$$

and so

$$\sum_{l=1}^L \|x_l\|_{\Omega_l} \geq \sum_{l:A \in \Omega_l} \|x_l\|_{\Omega_l} \geq \sum_{l:A \in \Omega_l} \frac{1}{2} |x_l(A)| \geq \frac{1}{2} |x(A)|,$$

then $\|x\| \geq \frac{1}{2} |x(A)| > 0$. □

Given a linear continuous functional $f : X \rightarrow \mathbb{R}$, define the game G_f as follows

$$G_f(A) = f(e_A)$$

for all $A \in \mathcal{C}$.

Theorem 5 *Let X' be the norm dual of $(X, \|\cdot\|)$. The operator*

$$\begin{aligned} G : X' &\rightarrow BV \\ f &\mapsto G_f \end{aligned}$$

is an isometric isomorphism from X' onto BV . Moreover, G is a weak-vague homeomorphism.*

Notice that, together with the Alaoglu Theorem (see, e.g., [3] p. 70), the above result immediately yields Theorem 1, that is, the compactness of the unit ball $U(BV)$ in the vague topology.

Proof. We first show that if $\Omega = \{S_i\}_{i=0}^n$ is a chain in \mathcal{C} then

$$\sum_{k=1}^n |G_f(S_k) - G_f(S_{k-1})| \leq \|f\|,$$

which implies that $G_f \in BV$ and $\|G_f\| \leq \|f\|$.

Define $x \in X_\Omega$ by

$$\begin{aligned}
x(S_n) &= \operatorname{sgn}(f(e_{S_n}) - f(e_{S_{n-1}})), \\
x(S_n) + x(S_{n-1}) &= \operatorname{sgn}(f(e_{S_{n-1}}) - f(e_{S_{n-2}})), \\
&\dots \\
x(S_n) + x(S_{n-1}) + \dots + x(S_1) &= \operatorname{sgn}(f(e_{S_1}) - f(e_{S_0})), \\
x(S_0) &= 0,
\end{aligned}$$

Obviously $\|x\|_\Omega \leq 1$, so that $\|x\| \leq 1$. Thus

$$\begin{aligned}
\|f\| &\geq f(x) = \sum_{j=0}^n f(e_{S_j}) x(S_j) = \\
&= f(e_{S_0}) \sum_{k=0}^n x(S_k) + \sum_{j=1}^n \left[(f(e_{S_j}) - f(e_{S_{j-1}})) \sum_{k=j}^n x(S_k) \right] = \\
&= \sum_{j=1}^n [(f(e_{S_j}) - f(e_{S_{j-1}})) \operatorname{sgn}(f(e_{S_j}) - f(e_{S_{j-1}}))] = \\
&= \sum_{j=1}^n |f(e_{S_j}) - f(e_{S_{j-1}})| = \sum_{j=1}^n |G_f(S_j) - G_f(S_{j-1})|.
\end{aligned}$$

Then G is well defined and obviously linear and injective.

Given $u \in BV$, we can define f_u on X by

$$f_u(x) = \sum_{A \in \mathcal{C}} u(A) x(A),$$

for all $x \in X$. Trivially, f_u is linear.

If x depends on $\Omega = \{S_j\}_{j=0}^n$, then

$$\begin{aligned}
f_u(x) &= \sum_{j=0}^n u(S_j) x(S_j) = \\
&= u(S_0) \sum_{k=0}^n x(S_k) + \sum_{j=1}^n \left[(u(S_j) - u(S_{j-1})) \sum_{k=j}^n x(S_k) \right] = \\
&= \sum_{j=1}^n \left[(u(S_j) - u(S_{j-1})) \sum_{k=j}^n x(S_k) \right] \leq \\
&\leq \sum_{j=1}^n \left[|u(S_j) - u(S_{j-1})| \left| \sum_{k=j}^n x(S_k) \right| \right] \leq \\
&\leq \sum_{j=1}^n [|u(S_j) - u(S_{j-1})| \|x\|_\Omega] \leq \\
&\leq \|x\|_\Omega \sum_{j=1}^n |u(S_j) - u(S_{j-1})| \leq \|x\|_\Omega \|u\|.
\end{aligned}$$

If $x = \sum_{l=1}^L x_l$ with $x_l \in X_{\Omega_l}$ for all $l = 1, 2, \dots, L$, then

$$f_u(x) = \sum_{l=1}^L f_u(x_l) \leq \sum_{l=1}^L \|u\| \|x_l\|_{\Omega_l} \leq \|u\| \sum_{l=1}^L \|x_l\|_{\Omega_l},$$

and so

$$\begin{aligned} f_u(x) &\leq \inf \left\{ \|u\| \sum_{l=1}^L \|x_l\|_{\Omega_l} : x = \sum_{l=1}^L x_l, x_l \in X_{\Omega_l} \right\} = \\ &= \|u\| \|x\|. \end{aligned}$$

We conclude that $f_u \in X'$, $G_{(f_u)} = u$ and G is onto. For all $u \in BV$, $f_u = G_u^{-1}$ and $\|G_u^{-1}\| = \|f_u\| \leq \|u\|$. Therefore, for all $f \in X'$, $\|f\| = \left\| G_{(G_f)}^{-1} \right\| \leq \|G_f\|$ and G is an isometry.

Finally, let $\{f^a\}$ be a net in X' . We have that $f^a \xrightarrow{w^*} f$ iff $f^a(x) \rightarrow f(x)$ for all $x \in X$ iff $f^a(e_A) \rightarrow f(e_A)$ for all $A \in \mathcal{C}$ iff $G_{f^a}(A) \rightarrow G_f(A)$ for all $A \in \mathcal{C}$ iff $G_{f^a} \xrightarrow{\mathcal{C}} G_f$. Hence, G is a weak*-vague homeomorphism. \square

We conclude the section by observing that Theorem 5 corrects Theorem 1.1 in [5]: it can be shown with a counterexample that the norm used there does not lead to an isometry.

5 Summing up

In sections 3 and 4 we have given two independent proofs that BV is the norm dual of X , endowed with the norms $\|\cdot\|_{BV'}$ and $\|\cdot\|$, respectively. The two approaches are indeed one the mirror image of the other. In fact:

Proposition 6 *The norm $\|\cdot\|_{BV'}$ on X coincides with the norm $\|\cdot\|$ on X . Moreover, $J = G^{-1}$.*

Proof. Let $(X', \|\cdot\|')$ be the norm dual of $(X, \|\cdot\|)$. We have, for all x ,

$$\begin{aligned} \|x\|_{BV'} &= \sup \left\{ \left| \sum_{A \in \mathcal{C}} u(A) x(A) \right| : u \in BV, \|u\| = 1 \right\} = \\ &= \sup \{ |f_u(x)| : u \in BV, \|u\| = 1 \} = \\ &= \sup \{ |f(x)| : f \in X', \|f\|' = 1 \} = \\ &= \|x\|. \end{aligned}$$

In particular the norm dual of $(X, \|\cdot\|_{BV'})$ and the norm dual of $(X, \|\cdot\|)$ are the same space X' . For all $u \in BV$ and $x \in X$,

$$J_u(x) = \langle u, x \rangle = \sum_{A \in \mathcal{C}} u(A) x(A) = f_u(x) = G_u^{-1}(x),$$

that is $J_u = G_u^{-1}$ for all $u \in BV$ and $J = G^{-1}$. \square

References

- [1] R.J. Aumann and L.S. Shapley, “Values of Non-Atomic Games”, Princeton University Press, Princeton NJ, 1974.
- [2] N. Dunford and J.T. Schwartz, “Linear Operators”, Interscience, New York, 1958.
- [3] R.B. Holmes, “Geometric Functional Analysis and its Applications”, Springer-Verlag, New York, 1975.
- [4] M. Marinacci, Decomposition and Representation of Coalitional Games, *Math. Oper. Res.* 21 (1996), 1000-1014.
- [5] W.H. Ruckle, Projection in Certain Spaces of Set Functions, *Math. Oper. Res.* 7 (1982), 314-318.

INTERNATIONAL CENTRE FOR ECONOMIC RESEARCH
APPLIED MATHEMATICS WORKING PAPER SERIES

1. Luigi Montrucchio and Fabio Privileggi, "On Fragility of Bubbles in Equilibrium Asset Pricing Models of Lucas-Type," *Journal of Economic Theory*, forthcoming (ICER WP 2001/5).
2. Massimo Marinacci, "Probabilistic Sophistication and Multiple Priors," *Econometrica*, forthcoming (ICER WP 2001/8).
3. Massimo Marinacci and Luigi Montrucchio, "Subcalculus for Set Functions and Cores of TU Games," April 2001 (ICER WP 2001/9).
4. Juan Dubra, Fabio Maccheroni, and Efe Ok, "Expected Utility Theory without the Completeness Axiom," April 2001 (ICER WP 2001/11).
5. Adriana Castaldo and Massimo Marinacci, "Random Correspondences as Bundles of Random Variables," April 2001 (ICER WP 2001/12).
6. Paolo Ghirardato, Fabio Maccheroni, Massimo Marinacci, and Marciano Siniscalchi, "A Subjective Spin on Roulette Wheels," July 2001 (ICER WP 2001/17).
7. Domenico Menicucci, "Optimal Two-Object Auctions with Synergies," July 2001 (ICER WP 2001/18).
8. Paolo Ghirardato and Massimo Marinacci, "Risk, Ambiguity, and the Separation of Tastes and Beliefs," *Mathematics of Operations Research*, forthcoming (ICER WP 2001/21).
9. Andrea Roncoroni, "Change of Numeraire for Affine Arbitrage Pricing Models Driven By Multifactor Market Point Processes," September 2001 (ICER WP 2001/22).
10. Maitreesh Ghatak, Massimo Morelli, and Tomas Sjöström, "Credit rationing, wealth inequality, and allocation of talent", September 2001 (ICER WP 2001/23).
11. Fabio Maccheroni and William H. Ruckle, "BV as a Dual Space," *Rendiconti del Seminario Matematico dell'Università di Padova*, forthcoming (ICER WP 2001/29).
12. Fabio Maccheroni, "Yaari Dual Theory without the Completeness Axiom," October 2001 (ICER WP 2001/30).